

# REHAB IN REVIEW

WWW.REHABINREVIEW.COM

TM

Volume 22 Number 6

Published by Physicians  
In Physical Medicine and Rehabilitation

June 5, 2014

## BODY MASS INDEX AND STROKE REHABILITATION

While obesity is associated with adverse health consequences, including comorbidities associated with stroke, the relationship between body mass index (BMI) and recovery following stroke is not well understood. This study assessed the relationship between BMI and the patient's progress in an acute rehabilitation hospital.

This retrospective study included all patients admitted to the stroke unit of a freestanding rehabilitation hospital between January of 2000 and April of 2006. All subjects underwent measurement of height and weight on admission, with BMI calculated. The participants were separated into four BMI categories, including underweight ( $<18.5\text{kg/m}^2$ ), normal weight ( $18.51\text{--}25\text{ kg/m}^2$ ), overweight ( $25.1\text{--}30\text{ kg/m}^2$ ) or obese ( $>30\text{ kg/m}^2$ ). FIM scores were obtained upon admission and throughout the rehabilitation stay. The primary outcome measure was FIM efficiency, as defined by the change in FIM score/days.

During the study period, 819 patients underwent stroke rehabilitation. Of these, five percent were underweight, 42% were normal weight, 32% were overweight and 20% were obese. The overweight group had the highest FIM efficiency ( $p=0.05$  as compared with the obese group). After adjusting for age and gender, FIM efficiency-motor did not differ by weight category, while FIM efficiency-cognitive demonstrated better progress for the overweight group ( $p=0.05$ ).

**Conclusion:** This study of patients admitted to an inpatient rehabilitation hospital found that overweight patients have better functional progress than do those in other weight categories.

Burke, D., et al. Effect of Body Mass Index on Stroke Rehabilitation. *Arch*

*Phys Med Rehab.* 2014, June; 95(6): 1055-1059.

## AMBULANCE-BASED THROMBOLYSIS FOR ISCHEMIC STROKE

Tissue plasminogen activator (tPA) is the treatment of choice for patients with ischemic stroke, after exclusion of intracerebral hemorrhage. However, only a minority of stroke patients receive tPA, due in part to a delay between the onset of symptoms and treatment. This study was designed to determine whether beginning tPA in the ambulance may improve outcome.

The Prehospital Acute Neurological Treatment and Optimization of Medical Care in Stroke Study (PHANTOM-S) is a randomized, open label, clinical trial, conducted in Berlin, Germany. Intervention involved an in-ambulance Stroke Emergency Mobile (STEMO), equipped with a CT scanner, a laboratory and a specialized pre-hospital stroke team. For this study, weeks were allocated to include either STEMO services or routine care. During the weeks with STEMO, thrombolysis was started en route to the hospital if ischemic stroke was confirmed and contraindications excluded. The primary outcome measure was duration from alarm to thrombolytic treatment. Secondary outcomes included in-hospital mortality and tPA rates.

During the 21-month study period, data from 6,182 patients were included. The rate of tPA treatment was 32.6% during the weeks with STEMO deployment and 21% during control weeks ( $p<0.001$ ). Treatment began within 90 minutes from onset in 58% of the patients after STEMO deployment and 37% during control weeks ( $p<0.001$ ). Compared with control weeks, STEMO resulted in a reduction of 15 minutes in time to treatment. However, no significant

difference was seen between groups in the number of in hospital deaths. STEMO deployment resulted in no increase in risk for intracerebral hemorrhage or seven day mortality.

**Conclusion:** This study of patients with acute ischemic stroke found that ambulance-based thrombolysis can reduce the time to tissue plasminogen treatment, without increasing adverse effects.

Ebinger, M., et al. Effect of the Use of Ambulance-Based Thrombolysis on Time to Thrombolysis in Acute Ischemic Stroke: A Randomized, Clinical Trial. *JAMA.* 2014, April 23/30; 311(16): 1622-1631.

## MESENCHYMAL STEM CELLS FOR OSTEOARTHRITIC KNEES

Cartilage regeneration procedures have become the focus of increased interest due to their potential to provide pain relief and alter the progression of osteoarthritis (OA). While mesenchymal stem cells (MSCs) can be derived from a number of tissues, some have suggested that those derived from adipose have the highest chondrogenic potential. This study investigated outcomes of adipose derived MSC implantation in patients with OA of the knees.

This retrospective review included the records of 56 consecutive patients treated with MSC implantation for cartilage regeneration. All had isolated, full-thickness articular cartilage lesions in OA knees. The mean size of the cartilage lesions was  $5.4\text{ cm}^2$ . The subjects originally had MSCs aspirated from adipose tissue from their buttock fat pad, with the stem cells introduced to the cartilage lesion under arthroscopic guidance. The knees were then immobilized for two weeks, with weight bearing initiated at two weeks, progressing to full weight-bearing at four weeks. Clinical

### **Editor-in-Chief**

David T. Burke, M.D., M.A.  
Emory University, Atlanta, GA

### **Executive Editor**

Randolph L. Roig, M.D.  
Emory University, Atlanta, GA

### **Copy Editor**

Roberta Alysoun Bell, Ph.D.  
Emory University, Atlanta, GA

### **Contributing Editors**

\*Erika Moody, M.D.  
BCM/UT Alliance, Houston, TX

\*Jennifer Knowlton, M.D.  
East Carolina University, Greenville, NC

\*Bamidele Oyebamiji Adeyemo, M.D.  
Emory University, Atlanta, GA

\*Mikhail Zhukalin, D.O.  
Anna Cruz, M.D.  
Chukwuemeka Ibekwe, M.D.  
Mitchel Leavitt, M.D.  
Cleo D. Stafford II, M.D., M.S.  
Donnie Staggs, M.D.  
Christopher Williams, M.D.  
Emory University, Atlanta, GA

\*Cyrus Kao, M.D.  
\*Rajat Mathur, M.D.  
\*Pablo Vazquez, M.D.  
Georgetown/MedStar Nat. Rehab, Wash., DC

\*Alexander Drakh, D.O.  
Casey Murphy, D.O.  
Eric Sterne, M.D.  
Grant Stone, D.O.  
LSU Medical Center, New Orleans, LA

\*Bryndon Hatch, M.D.  
\*Katie Nanos, M.D.  
Mayo Clinic, Rochester, MN

\*Eric Leung, M.D.  
Mount Sinai Med. Ctr., New York, NY

\*Arpit A. Patel, D.O.  
Burhan Haleem, D.O.  
Ashley Zakhary, M.D.  
Nassau University Med Center

\*Christina Marciniak, M.D.  
Mary Caldwell, D.O.  
Leda Ghannad, M.D.  
Nying Lam, M.D.  
Amy Matthews, M.D.  
Sangeeta Patel, M.D., MPH  
Nicole Rup, M.D.  
N.W.U./R.I.C., Chicago, IL

\*David Woznica, M.D.  
NYP, Columbia-Cornell, NY, NY

\*Samaira Khan, D.O.  
NYU/Rusk Inst. of Rehab Med, NY, NY

\*John Furrey  
Rush Univ. Medical Center, Chicago, IL

\*Leroy Lindsay, M.D.  
Michelle Didesch, M.D.  
Rutgers/Kessler Rehab, Newark, NJ

\*Catherine J. Yee, M.D.  
Karen Cruz, D.O.  
Zainab Najji, M.D.  
Holly Pajor, D.O.  
Schwab Rehab/U. of Chicago Hospitals, IL

\*Alan Vo, D.O.  
Sinai Hospital, UMD, Baltimore, MD

\*Anthony A. Cuneo, M.D., PhD  
Joyce Cheng, M.D.  
Adam Weidenhammer, M.D.  
Temple Univ./UPenn., Philadelphia, PA

\*Anupam Sinha, D.O.  
Th. Jefferson U./Rothman Inst., Phil., PA

\*Deep Garg, M.D.  
Lauren Lerner, M.D.  
Joshua Rothenberg, D.O.  
University of Miami, Miami, FL

\*Seth Swank, D.O.  
Univ. of Michigan, Ann Arbor, MI

\*Peter Hwang, M.D.  
University of Missouri, Columbia MO

\*Alexis Tingan, M.D.

outcomes were evaluated using the International Knee Documentation Committee (IKDC) score and Tegner activity scale. For those agreeing to a second look arthroscopic assessment, cartilage repair was assessed using International Cartilage Repair Society (ICRS) grading.

At final follow-up, the mean IKDC and Tegner activity scale scores had both improved significantly ( $p < 0.001$ ). The patients reported that satisfaction with the procedure was excellent in 63%, and good in 31%, of the cases. However, at second look arthroscopy, two of the 37 lesions (five percent) had grade I (normal), seven (19%) had grade II (near normal), 20 (54%) had grade III (abnormal) and eight (22%) had grade IV (severely abnormal) lesions. Those patients with lesions larger than 5.4 cm<sup>2</sup> had significantly worse outcomes than did those with smaller lesions.

**Conclusion:** This study of patients with osteoarthritis of the knee found that autologous mesenchymal stem cell implantation, derived from adipose, can produce significant functional gains.

Koh, Y., et al. Second Look Arthroscopic Evaluation of Cartilage Lesions after Mesenchymal Stem Cell Implantation in Osteoarthritic Knees. *Am J Sports Med.* 2014, doi:10.1177/0363546514529641

### **AUTOLOGOUS CHONDROCYTE IMPLANTATION AT THE PATELLA**

Cartilage defects are present in more than 50% of patients undergoing knee arthroscopy. The articular surface of the patella is affected in 11 to 36% of the cases. However, cartilage repair at the patella has demonstrated relatively poor results. This study assessed the effect of treatment with autologous chondrocyte implantation (ACI) for cartilage defects of the patella.

Patients presenting for repair of cartilage defects in the patella underwent chondrocyte harvesting, with chondrocytes expanded in cell culture. The cells were then implanted at the site of the defects. Postoperative rehabilitation included the use of a locked brace for ambulation, with weight bearing as tolerated for those with isolated ACI. The patients were followed prospectively, with questionnaires

used to assess function, activity level and satisfaction.

A total of 110 patients were followed post-surgery for an average of 90 months. The average size of the patella defect was 5.4 cm<sup>2</sup>. Nine were considered treatment failures and were revised. Overall, the Short Form -12 Physical Component Subscale demonstrated an increase from 38.6 to 44.1 ( $p = 0.001$ ). Significant improvements were also noted in the International Knee Documentation Committee (IKDC) scores ( $p < 0.0001$ ), the modified Cincinnati score ( $p < 0.0001$ ), the Knee Society knee score, and the Knee Society function score, ( $p < 0.0001$ ). The WOMAC score improved from 50.4 to 28.6, with 75% of patients exceeding a commonly accepted threshold for MCIDs. Of the 93 patients asked to rate their knee, 86% rated their knee function as good or excellent.

**Conclusion:** This study of patients with cartilage defects in the patella found that autologous chondrocyte implantation can produce significant and clinically meaningful improvements in pain and function.

Gomoll, A., et al. Autologous Chondrocyte Implantation in the Patella: A Multicenter Experience. *Am J Sports Med.* 2014, May; 42(5): 1074-1081.

### **STEM CELLS FOR ROTATOR CUFF HEALING**

Rotator cuff tears occur in more than 50% of individuals over the age of 60 years. However, approximately 50% of surgically repaired rotator cuffs do not heal properly. As stem cell therapies for tissue regeneration and repair represent a new avenue for investigation, this study was designed to determine whether healing after a rotator cuff repair can be accelerated through the use of autologous stem cells.

This animal study included 40 white male rabbits. All underwent bilateral subscapularis (SSC) rupture. After six weeks the rabbits were randomly divided into five groups. These included adipose-derived stem cells (ADSCs) plus repair, saline plus repair, ADSCs only, saline only and a control. Animals then underwent electromyographic, biomechanical and histologic analysis six weeks after the secondary procedures.

All tendons in the ADSC only and saline groups failed to heal. Biomechanically, the load to failure of the ADSC plus repair group was 87.02 N, and that of the saline plus repair group was 59.85 N ( $p=0.085$ ). Histologically, the mean proportion of fatty infiltration in the SSC muscles was lowest in the ADSC plus repair group ( $p<0.001$ ).

**Conclusion:** This study, using an animal model for rotator cuff tears, found that locally administered adipose derived stem cells may have a positive benefit on tendon-to-bone healing and fatty infiltration.

Oh, J., et al. 2013 Neer Award: Effect of the Adipose Derived Stem Cell for the Improvement of Fatty Degeneration and Rotator Cuff Healing in a Rabbit Model. *J Shoulder Elbow Surg.* 2014, April; 23(4): 445-456.

#### FACTORS PREDICTING ROTATOR CUFF RETEARS

After rotator cuff tears, the rate of retears is estimated to be between 11 and 57%. Factors that contribute to rotator cuff retears remain unclear. Thus, this study was designed to better understand risk factor for retears.

This retrospective study included 1,000 consecutive patients who had undergone a primary rotator cuff repair. Preoperatively, all patients completed a questionnaire grading the frequency of pain during activity and at rest, as well as the frequency of extreme shoulder pain and stiffness. Questions also focused on employment and sport related activity. All patients underwent a preoperative clinical examination. During surgery, measurements were made of the tear. Six months after surgery, all participants underwent ultrasound examination to determine the integrity of the surgical site.

At surgery, 561 full thickness and 393 partial thickness tears were noted. At follow-up, retears were seen in 174 patients, corresponding to an overall rate of 17%. Independent factors predicting rotator cuff retears, ranked by strength of predictive value, included anterior posterior tear lengths ( $p<0.001$ ), age at surgery ( $p<0.001$ ), tear size area ( $p<0.001$ ), mediolateral tear length ( $p<0.001$ ), tear thickness ( $p=0.022$ ) and operative time ( $p=0.036$ ). Using

these, the authors provide an equation for estimating the risk for retear.

**Conclusion:** This retrospective analysis of 1,000 consecutive rotator cuff retears found that initial tear dimensions and advanced age were significantly related to the risk of retear at six months.

Le, B., et al. Factors Predicting Rotator Cuff Retears. An Analysis of 1,000 Consecutive Rotator Cuff Repairs. *Am J Sports Med.* 2014, May; 42(5): 1134-1142.

#### NSAIDS AFTER ROTATOR CUFF TENDON REPAIR

While nonsteroidal anti-inflammatory drugs (NSAIDs) have been found to affect bone metabolism and fracture healing, little evidence exists concerning their effect on tendon healing. This animal study assessed the effect of a selective COX-2 inhibitor on tendon-bone healing.

Using an animal rotator cuff model, surgery was performed on 39 rats to detach and repair the supraspinatus tendon insertion sites. After surgery, 13 rats received daily injections of meloxicam for 10 days (group A), while 13 rats received injections of meloxicam starting at postoperative day 11 (group B). A third group of 13 rats received saline injections (group C). Tendon specimens of each group were used for gross inspection, histopathology evaluation and biomechanical testing. Biomechanical testing was completed at 21 days post-surgery, using parameters including load to failure, stiffness and displacement.

Maximal loads in group B were significantly lower than those for both the C and A groups. Stiffness was lower in group B than in group A ( $p=0.05$ ). Histological examination revealed less organized collagen at the tendon bone interface in group B than in the other two groups.

**Conclusion:** This animal study found that, after a rotator cuff repair, delayed (days 11 to 20) use of nonsteroidal anti-inflammatory drugs, during the proliferative phase of healing may negatively impact tendon strength and collagen formation.

Chechik, O., et al. Timing Matters: NSAIDs Interfere with the Late Proliferation Stage of the Repaired

Rotator Cuff Tendon Healing in Rats. *Arch Orthop Trauma Surg.* 2014, April; 134(4): 515-520.

#### ANGIOTENSIN II TYPE 2 RECEPTOR ANTAGONIST FOR POST-HERPETIC NEURALGIA

Findings from earlier studies have demonstrated that the Angiotensin II, type 2 receptor (AT<sub>2</sub>R) is expressed in human sensory neurons. As studies have demonstrated that EMA401, a highly selective AT<sub>2</sub>R antagonist, has analgesic properties, this study was designed to further determine the analgesic efficacy, tolerability, safety and kinetics of this compound for patients with post-herpetic neuralgia.

This multicenter, double-blind, placebo-controlled trial included participants from 29 sites in six countries. All patients had a diagnosis of post-herpetic neuralgia, and were randomized to receive either the study drug, at 100 mg, or a placebo, administered twice per day. After randomization, the subjects completed efficacy endpoint patient outcome measures, and were given a diary card to record their daily pain intensity and the time of study drug administration. All were assessed for efficacy, safety and pharmacokinetics. The primary endpoint was the change in mean pain intensity from baseline to the final week of dosing.

A total of 183 patients were assessed, with 92 receiving the study drugs and 91 the placebo. The changes in mean pain intensity between baseline and final weeks of treatment were 2.29 for the treatment group and 1.6 for the placebo group ( $p=0.0066$ ). The difference between the groups in mean pain intensity first reached significance at week three. No significant treatment related side effects were noted in the treatment group.

**Conclusion:** This study of patients with post-herpetic neuralgia found that EMA401, a highly selective angiotensin II type 2 receptor antagonist, significantly reduced the pain symptoms of this condition.

Rice, A., et al. EMA401, an Orally Administered, Highly Selective Angiotensin II Type 2 Receptor Antagonist, as a Novel Treatment for Post-Herpetic Neuralgia: A Randomized, Double-Blind, Placebo-

Controlled, Phase 2 Clinical Trial. **Lancet**. 2014, May 10-16: 1637-1647.

### MEMORY CONSOLIDATION WITH CAFFEINE

Caffeine has been shown to have positive effects on attention, processing speed and, in some cases, working memory. However, most studies have administered caffeine before learning, causing the effects on memory to be impossible to dissociate from other effects of the caffeine. This study assessed the effect of caffeine administered after a learning task.

This randomized, double-blind, placebo-controlled trial included caffeine naïve participants. A hippocampal memory-dependent task included two phases. During the incidental encoding phase, participants viewed pictures of objects and decided whether the picture depicted an indoor or an outdoor item. The subjects were then given either a placebo or 200 mg of caffeine. Twenty-four hours after the study session, the participants were evaluated for recognition performance, with researchers presenting items from the previous day (targets), new items (foils) and items that were similar but not identical to previously shown pictures (lures). For each image, they were instructed to decide whether the image was "old", "new" or "similar".

Participants who received caffeine were more likely to call lure items "similar", rather than "old", as compared to those who received placebo. No difference was found between groups in rates of target hits or foil rejections, suggesting that basic recognition was not altered. Calculating a lure discrimination index, as  $P$  ("similar"/lure) minus  $P$  ("similar"/foil) to correct for response bias, a significant difference was found between groups ( $p=0.05$ ). The authors note that this finding suggests that caffeine enhanced consolidation of the initial study session, such that discrimination during retrieval was improved. Reviewing multiple caffeine doses, it was determined that a dose of at least 200 mg was required to produce the enhancing effect of caffeine on consolidation of memory.

**Conclusion:** This study found that caffeine, administered after

information encoding, improved the patients' ability to consolidate memories.

Borota, D., et al. Post- Study Caffeine Administration Enhances Memory Consolidation in Humans. **Nature Neurosci**. 2014, February; 17(2): 201-203.

### ENERGY EXPENDITURE DURING RESISTIVE TRAINING ACTIVITY

To date, few studies have addressed the energy expenditure of resistance training. This study assessed the energy expenditure of resistance training using two separate methods.

Subjects included twelve healthy men, with a mean age of 23.6 years. The subjects completed each individual resistive training exercise for a maximum of 60 seconds or until fatigue degraded the proper form. After completing each exercise, the subject was rested. Traditional calorimetry was used to measure oxygen uptake continuously throughout the trial, including during the recovery period between exercises. The average energy expenditure was calculated using a traditional calculation as well as a recovery calculation.

Based upon the traditional calculation, the kilocalories-per-minute expenditure was 4.31 for push-ups, 4.09 for curl-ups, 5.28 for lunges and 4.03 for pull-ups. However, using the recovery calculation, the kilocalories-per-minute were 8.56 for push-ups, 7.29 for curl-ups, 9.33 for lunges and 9.95 for pull-ups. After converting the mean values for each exercise to METS, all were classified as being of moderate intensity when using the traditional method, while when using the recovery calculation, all but the curl-ups were classified as vigorous.

**Conclusion:** This study of energy expenditure during resistive training found that traditional measures of energy expenditure may underestimate energy consumed during these activities, with the authors suggesting that the recovery time method of estimation may be more accurate.

Vezina, J., et al. An Examination of the Differences between Two Methods of Estimating Energy Expenditure in Resistive Training

Activities. **J Strength Cond Res**. 2014, April; 28(4): 1026-1031.

### DONEPEZIL AND LIFE EXPECTANCY

A number of studies have demonstrated that cognitive impairment with advancing age is a negative predictor of survival. Drugs such as cholinesterase inhibitors have been used for symptomatic treatment of dementias, including Alzheimer's disease (AD). This study was designed to determine whether donepezil, a cholinesterase inhibitor, affects life expectancy.

This study included outpatients at a memory clinic seen between 1999 and 2012. All had undergone magnetic resonance imaging, with study entry criteria including dementia with a Clinical Dementia Rating of 1+, medical treatment for more than three months and follow-up until less than one year before death. The main outcome variable was the time between the onset of AD and death, compared by use of donepezil. A one-way analysis of variance was performed to analyze the effect of donepezil administration on life expectancy.

Of the 390 medical records reviewed, 275 patients diagnosed with dementia met the entry criteria. Of the 100 patients with AD, and AD with cerebrovascular disease, 52 received donepezil and 48 did not. Life expectancies after AD onset were 7.9 years in the donepezil group and 5.3 years in the non-donepezil group ( $p<0.001$ ). The Spearman correlation found a positive correlation between the period of donepezil use and life expectancy in AD ( $p<0.0001$ ).

**Conclusion:** This retrospective study of patients with Alzheimer's disease found that treatment with donepezil may increase life expectancy.

Meguro, K., et al. Donepezil and Life Expectancy in Alzheimer's Disease: A Retrospective Analysis In the Tajiri Project. **BMC Neurol**. 2014; 14: 83.

### TENNIS-RELATED INJURIES

Tennis is among the fastest growing sports in the United States, with a 46% increase in participants over the past decade. Most data

concerning tennis injuries are based upon elite or professional, rather than amateur, player data. This study was designed to determine patterns of injury among all tennis players seen for injury in emergency departments in the United States.

Data were obtained from the National Electronic Injury Surveillance System (NEISS), with data collected from 100 hospitals, representing a stratified probability sample of 5,400 hospitals. The data were reviewed for mechanism of injury and reviewed by age categories. Injury diagnoses and locations were identified.

From 1990 through 2011, 492,002 patients were treated in United States emergency departments for tennis-related injuries. The number of tennis related injuries decrease by 41.4% during the years 1990 to 2011. Children five to eighteen years of age accounted for 29.2% of those injuries, and patients 19 to 45 years accounted for 27.3% of the cases. Most patients were treated and released (96.2%). Among the patients admitted to a hospital, 65.6% were 56 years of age or older.

The most commonly injured body regions were lower extremities, representing 42.2% of the injuries, with injuries to the ankles representing 47.2% of lower extremity injuries. The wrist represented 34.8% of upper extremity injuries. Strains and sprains were the modal injury, at 44.1%, followed by fractures, at 14.6%, and soft tissue injuries, at 13.9%.

**Conclusion:** This study demonstrates that tennis-related injuries have decreased over the past 20 years. More than 40% of all injuries involved the lower extremities.

Gaw, C., et al. Tennis Related Injuries Treated in the United States Emergency Departments, 1990 to 2011. *Clin J Sport Med*. 2014, May; 24(3): 226-232.

### HAND OSTEOARTHRITIS AND FENOFIBRATE

Fibrates are agonists of the peroxisome proliferator activated receptor alpha. These medications have long been used for dyslipidemia and have also demonstrated benefits for reducing systemic and synovial inflammation, inhibit chondrocyte inflammatory response and exert anti-

nociceptive properties. This study was designed to determine the effects of fenofibrate for the treatment of patients with erosive hand osteoarthritis (EHOA).

This single center non-controlled open label study included 15 patients over the age of 45 years with a diagnosis of HOA. The patients received treatment with 145 mg of fenofibrate per day for 12 weeks. Clinical evaluations were made at baseline and at weeks four, eight, and 12. These included assessment of disease activity, pain, and laboratory tests of inflammatory markers. Response to treatment was assessed in accordance with Outcome Measures in Rheumatology Clinical Trials and Osteoarthritis Research Society criteria.

Fourteen women were enrolled in the study. Fenofibrate treatment resulted in significant decreases in pain scores, total joint count, the duration of morning stiffness, as well as the patient's assessment of disease activity and global health. An improvement of hand function was noted by improvements in the Cochin index. Significant reductions were noted in levels of ESR, while no change was noted in several other inflammatory markers.

**Conclusion:** This pilot, non-controlled study of patients with erosive hand osteoarthritis found that fenofibrate, an agonist of the peroxisome proliferator activated receptor alpha, may be helpful for reducing pain, and improving hand function.

Shirinsky, I et al. Treatment of Erosive Osteoarthritis with Peroxisome Proliferator – Activated Receptor Alpha Agonist Fenofibrate: A Pilot Study. *Rheum International* 2014, May. 34(5):613-616.

### EXERCISE INTERVENTIONS TO PREVENT SPORTS INJURY

As management of sports injuries is difficult and time-consuming, prevention of these injuries is of particular interest. This meta-analysis was designed to better understand the effect of physical activity exercises as a means to prevent injuries.

Data were reviewed from several databases, searched through January of 2013. Those studies included focused on primary prevention,

subjects free of injury at inclusion, sports/physical activity injuries, randomized controlled trials, appropriate controls and publication in peer reviewed journals.

Twenty-five trials were selected for inclusion, including data from 26,610 individuals with 3,464 injuries. Injury prevention strategies were assessed. As a group, the relative risk of injury, using the prevention strategies was 0.632, with significant findings for both adults and adolescents. The pooled effect of strength training, including four studies, resulted in a relative risk of 0.315. The pooled effect of proprioceptive training resulted in a relative risk of 0.55. No significant effect was found for stretching.

**Conclusion:** This review and meta-analysis of injury prevention programs found that, while stretching has no beneficial effect, multiple exposure programs, proprioceptive training and strength training are effective strategies. Strength training was found to reduce injury risk to less than one third.

Lauersen, J., et al. The Effectiveness of Exercise Interventions to Prevent Sports Injuries: A Systematic Review and Meta-Analysis of Randomized, Controlled Trials. *Br J Sports Med*. 2014, June; 48(11): 871-877.

### STRETCHING TECHNIQUES AND VERTICAL JUMPING PERFORMANCE

Many coaches believe that static stretching before competition improves performance, prevents injuries and increases flexibility. Recent studies have suggested, in fact, that static stretching may negatively impact athletic performance, and that the effect of stretching on performance depends upon the type of stretching. This study evaluated three different stretching techniques with regard to their effect on vertical jumping performance.

One hundred, physically active, male athletes volunteered for participation in this study. All subjects underwent pre-jump performance assessments, including flexibility measurement. The volunteers were divided into three groups, according to flexibility and pre-jump performances. After warm-up, the participants were asked to perform

one of three stretching techniques. Those techniques included ballistic stretching (BS), proprioceptive neuromuscular facilitation stretching plus ballistic stretching (PNF plus BS) and proprioceptive neuromuscular facilitation stretching plus static stretching (PNF plus SS). Each stretching technique was applied bilaterally, with four sets of each performed. The subjects were then assessed by countermovement jump, and were asked to jump as high as possible while extending the legs. The jump was repeated three times.

For the entire group, BS increased vertical jump performance, and also improved the performance of those in the low average flexibility and poor pre-jumping performance groups. PNF plus BS improved performance among those with high flexibility. Proprioceptive neuromuscular facilitation plus SS decreased performance in the whole group, as well as among those in the high flexibility, moderately high pre-jumping performance groups.

**Conclusion:** This study, comparing pre-performance stretching techniques, found that combining proprioceptive neuromuscular facilitation with static stretching can degrade jumping performance. The results suggested that ballistic stretching is more suitable, resulting in enhanced performance.

Kirmizigil, B., et al. Effects of Three Different Stretching Techniques on Vertical Jumping Performance. *J Strength Cond Res.* 2014, May; 28 (5): 1263-1271.

#### OSTEOARTHRITIS AFTER ANTERIOR CRUCIATE LIGAMENT RECONSTRUCTION

Estimates of the prevalence of osteoarthritis (OA) following anterior cruciate ligament (ACL) reconstruction range from 10% to 90%. Some have reported that meniscus injuries requiring repair increase this risk. However, randomized trials are sparse. This study was designed to better understand the prevalence of radiological OA after ACL reconstruction.

This randomized, controlled trial included 164 patients, treated between 1995 and 1997. Each had a traumatic ACL injury. The participants

were randomized to receive either a bone tendon bone graft or a semitendinosus graft. Fourteen years after reconstruction, 135 were contacted and agreed to participate in this follow-up study. All subjects underwent radiological examinations of both knees to determine the grade of OA, and all also underwent knee function assessment using a visual analogue scale and the Knee Injury and Osteoarthritis Outcomes Score (KOOS). Osteoarthritis was defined as a Kellgren-Lawrence grade 2 or greater.

Osteoarthritis was more frequent in the reconstructed knee than in the contralateral knee. OA of the medial compartment was most frequent, occurring in 57% of OA cases in the reconstructed knee, and in 18% in the contralateral knee ( $p<0.001$ ). Univariable analysis revealed that the risk of OA did not differ between those with, versus without medial meniscal injury. Patients with a resected medial meniscus were at increased risk relative to those with a repaired medial meniscus ( $p=0.03$ ).

**Conclusion:** This prospective study found a three-fold increase in osteoarthritis after an anterior cruciate ligament injury, treated with reconstruction, as compared to the contralateral uninjured knee.

Barenius, B., et al. Increased Risk of Osteoarthritis after Anterior Cruciate Ligament Reconstruction. A 14-Year, Follow-up Study of a Randomized, Controlled Trial. *Am J Sports Med.* 2014, May; 42(5): 1049-1057.

#### PSOAS TENDINOPATHY INTERVENTIONS

Psoas tendinopathy, or internal snapping hip syndrome, occurs predominantly as an overuse injury. Pain is thought to originate from the friction of the iliopsoas tendon sheath over the iliopectineal ridge or the iliacus tendon. Operative interventions include step cutting the iliopsoas and releasing the iliopsoas tendon. This study assessed the efficacy of surgery as well as tendon and bursa hip injections over time.

Patients with primary psoas tendinopathy were seen by a hip specialist and referred for musculoskeletal ultrasound and plain film radiographs. If snapping was visualized, triamcinolone and anesthetic were placed in the tendon

sheath and bursa. Six to eight weeks post-injection, the subjects were reassessed for psoas related pain. Those with no pain relief were removed from the study. Those with temporary benefit were offered the choice of further injections or surgical intervention. All were administered questionnaires, the nonarthritic hip score (NAHS) and a study questionnaire, which queried the patient concerning hip function activity and pain over the previous 48 hours.

From 1997 to 2011, 37 hips were identified with primary iliopsoas tendinitis. Of those, 13 hips were injected, with no pain improvement. Twenty-six hips, (70%) were treated surgically with psoas tenotomy, and 11 (30%) were treated by steroid injection alone. The average follow-up period for those patients treated by surgical release was 49 months, while the average for those treated with steroid injection was 52 months. The average NAHS score after surgery and injection were 68 and 86, respectively ( $p<0.05$ ). Improvements in pain were similar between the surgery and injection groups, though neither group noted increased activity.

**Conclusion:** This study of patients with primary psoas tendinopathy found that local steroid injections can provide long-term pain relief. For patients with only temporary relief from injections, psoas tenotomy can provide good long term pain relief.

Garala, K., et al. Medium Term and Long-Term Outcomes of Interventions for Primary Psoas Tendinopathy. *Clin J of Sports Med.* 2014, May; 24(3): 205-210.

#### FRACTURE AFTER ALENDRONATE DISCONTINUATION

The Fracture Intervention Trial Long-Term Extension (FLEX) trial demonstrated that, among older postmenopausal women who have used alendronate for five years, those who discontinue treatment have similar fracture rates to those who receive an additional five years of treatment. However, the rate of symptomatic vertebral fractures is lower among those who continue therapy. This post-hoc analysis was designed to test methods of

predicting fractures among women who have discontinued alendronate therapy.

Data were obtained from the FLEX trial, which randomized postmenopausal women to receive either alendronate sodium or placebo for five years. Bone mineral density (BMD) was measured at the hip and lumbar spine at baseline, and at the 36-month follow-up visit. In addition, bone turnover biochemical markers were assessed at baseline and at 12-month and 36-month visits. The primary outcome variable was the incidence of clinical fractures, defined as either new, non-spine fracture or new clinical vertebral fracture, during the FLEX follow-up period.

Of the 437 participants randomized to receive placebo, 22% experienced one or more clinical fractures during follow-up. Older age and lower hip BMD at the time of medication discontinuation were strong predictors of fracture risk after discontinuation. However, neither the one-year change in hip bone mineral density, nor the one- or three-year change in biochemical markers of bone turnover, were associated with the risk of fracture after discontinuation.

**Conclusion:** This study of postmenopausal women, all treated with alendronate therapy for five years, found that age and bone mineral density at the time of discontinuation of medication were useful for predicting clinical fractures during the subsequent five years.

Bauer, D., et al. Fracture Prediction after Discontinuation of Four to Five Years of Alendronate Therapy. The Flex Study. *JAMA Internal Med.* 2014; doi:10.1001/jamainternmed.2014.1232

### LOCAL COOLING FOR TRAUMATIC SPINAL CORD INJURY

In severe spinal cord injury (SCI), the primary injury forces destroy or damage neural tissue, with a secondary autodestructive process leading to infarction of cord tissues. This study evaluated whether those secondary processes can be arrested through a combination of steroids, decompressive surgery and localized cooling.

This prospective case series enrolled 20 patients over 10 years.

The subjects were between 16 and 65 years of age, and all were admitted with neurologically complete, ASIA A SCI. Patients received six milligrams of dexamethasone every six hours for 11 days tapering off until day 18. Surgical decompression with fusion was performed. Normal saline at 3<sup>0</sup> C was applied against the dura for four hours, cooling the dura to 6<sup>0</sup> C. The target time from injury to the initiation of cooling was eight hours. Following intervention, the participants were followed at intervals ranging from 14 to 153 months.

At final follow-up, an average of 4.9 years after injury, 35% of the patients remained categorized as ASIA A, 30% improved to ASIA B, 25% improved to ASIA C and 10% improved to ASIA D. For comparison authors cited data collected from the Model Spinal Cord Injury System. These data reveal a 3.5% chance of status improving to ASIA Grade B and a 1.05% chance of improving to ASIA Grade C or D within 5 years of a complete injury.

**Conclusion:** This small study of patients admitted with neurologically complete spinal cord injury found that a combined treatment of steroids, surgical decompression and local cooling resulted in better neurologic recovery than expected, as compared to patients receiving traditional treatment.

Hansebout, R., et al. Local Cooling for Traumatic Spinal Cord Injury: Outcomes in 20 Patients and Review of the Literature. *J Neurosurg Spine.* 2014, May; 20(5): 550-561.

### INTRACRANIAL ATHEROSCLEROSIS AND RISK OF STROKE

The lifetime risk of stroke is estimated to be at least one in six, with the burden in the upcoming decades expected to increase. Several locations of vascular disease are considered to be relevant to stroke etiology. Little data exist concerning the burden of stroke attributable to intracranial atherosclerosis in the white population. This study was designed to estimate the total burden of stroke attributable to intracranial carotid artery calcification (ICAC) in white individuals.

This study was based on the Rotterdam study, a prospective, population-based trial investigating the determinants of chronic diseases in the elderly. For this study, visits between September 1, 2003, and February of 2006 were used as a baseline. All subjects underwent non-enhanced CT scan in order to identify the volume of the ICAC. The patients were then monitored for six years for the development of stroke. Data from the population of white descent was used for analysis.

During 14,055 person-years of follow-up, 91 participants suffered a stroke, among which 74 were ischemic. Intracranial carotid artery calcification played a role in up to 75% of all strokes, while aortic arch and extracranial artery calcification contributed to 45% and 25% respectively. Further, large ICAC volume was associated with a higher risk of stroke, independent of other cardiovascular risk factors.

**Conclusion:** This study suggests that intracranial atherosclerosis is a major risk factor for the development of stroke in the general white population.

Bos, D., et al. Intracranial Carotid Artery Atherosclerosis and the Risk of Stroke in Whites: The Rotterdam Study. *JAMA Neuro.* 2014, April; 71 (4): 405-411.

### PULMONARY EMBOLCTOMY IN ELDERLY PATIENTS

Previous studies have demonstrated that age is an important factor among patients undergoing embolectomy after a pulmonary embolism. This study further explored the impact of age on case fatality rate.

Data were gathered from the Nationwide Inpatient Sample, which comprises data from 1,000 hospitals. Data were reviewed to identify patients with pulmonary embolism who underwent pulmonary embolectomy between 1999 and 2008. Those subjects were stratified according to whether they were medically stable or unstable. The data were then reviewed to determine the impact of age on fatality rate.

Between 1999 and 2008, 2,685, stable patients ages 21 years or older underwent pulmonary embolectomy. A sharp decline was seen in the proportion of stable patients



(Continued from page 2)

Lauren Lerner, M.D.  
Joshua Rothenberg, D.O.  
University of Miami, Miami, FL

\*Seth Swank, D.O.  
University of Michigan, Ann Arbor, MI

\*Peter Hwang, M.D.  
University of Missouri, Columbia MO

\*Alexis Tingan, M.D.  
UPenn Health System, Philadelphia, PA

\*Thiru Annaswamy, M.D.  
Trixy S  
UTSW Medical Center, Dallas TX

\*Rachel Hallmark, M.D., Ph.D.  
UVA, Charlottesville, VA

\*Ryan Solinsky, M.D.  
Debbie Tan, M.D.  
University of Washington, Seattle, WA

\*Bonnie Weigert, M.D.  
University of Wisconsin, Madison, WI

\*William Robbins, M.D.  
VCU, Richmond, VA

\*Adam J. Schulte, M.D.  
Shannon Dietzmann, M.D.  
Prateek Grover, M.D.  
Sri Patchala, M.D.  
Washington University, St. Louis, MO

#### **Executive Editor Emeritus**

Donald F. Langenbeck, Jr., M.D.

#### **Subscription Manager**

Michael P. Burke, M.S.

**\*Regional Managing Editors have attested that they have no financial conflict of interest when choosing articles that appear in Rehab in Review.**

undergoing embolectomy with increasing age, from 0.13% among those 71 to 80 years of age to 0.04% among those over 80 years of age ( $p < 0.001$ ). The case fatality rate with pulmonary embolectomy in stable patients increased with age beginning at age 51 to 60 years (17.4%), increasing to 46.2% among patients greater than 80 years of age ( $p < 0.0001$ ).

During the same time period, 925 unstable patients with pulmonary embolism underwent embolectomy. The rate of pulmonary embolectomy in that group was 10-fold greater than that among stable patients. However, the case fatality rate was not significantly related to age in the unstable group.

**Conclusion:** This study found that the all-cause case fatality rate among stable patients undergoing pulmonary embolectomy increases at age 51 to 60 years, while the fatality rate for unstable patients was not related to age.

Stein, P., et al. Pulmonary Embolectomy in Elderly Patients. **Am J Med.** 2014, April; 127(4): 348-350.

*Rehab in Review (RIR)* is produced monthly by physicians in the field of Physical Medicine and Rehabilitation (PM&R), with the cooperation and assistance of Emory University School of Medicine, Department of Rehabilitation Medicine. The summaries appearing in this publication are intended as an aid in reviewing the broad base of literature relevant to this field. These summaries are not intended for use as the sole basis for clinical treatment, or as a substitute for the reading of the original research.

The Emory University School of Medicine designates this journal based activity for a maximum of 3 AMA PRA Category 1 Credits™. Physicians should only claim credit commensurate with the extent of their participation in the activity. The Emory University School of Medicine is accredited by the ACCME to provide continuing medical education for physicians.

*RIR* is affiliated with the Association of Academic Physiatrists, the World Health Organization, and the Chinese and Indian Societies of PM&R and endorsed by the International Society of Physical and Rehabilitation Medicine.

Private subscriptions are available by email at [rehabinreview@aol.com](mailto:rehabinreview@aol.com) or by fax or phone at (800) 850-7388.

ISSN # 1081-1303

[www.rehabinreview.com](http://www.rehabinreview.com)



## **REHAB IN REVIEW**

Produced by the Department of  
Rehabilitation Medicine, Emory  
University School of Medicine



**EMORY**  
UNIVERSITY  
SCHOOL OF  
MEDICINE

Department of  
Rehabilitation  
Medicine

*Expanding the frontier of rehabilitation sciences in research, teaching, and patient care*